Prolonged Grief Disorder (PG-13-Revised)

Holly G. Prigerson, Ph.D., Jiehui Xu, M.S., Paul K. Maciejewski, Ph.D.

Q1. have you lost someone significant to you? (•) Yes / ( ) No

Q2. how many months has it been since your significant other died? _____ Months.

For each item below, please indicate how you currently feel?

Since the death, or as a result of the death…

		Not at all (0)	Slightly (1)	Somewhat (2)	Quite a bit (3)	Overwhelmingly (4)
Q3	**Do you feel yourself longing for the person who died?**					
Q4	**Do you have trouble doing things you'd usually do because you are thinking so much about the person who died?**					
Q5	**Do you feel confused about your role in life or feel like you don't know who you are anymore?**					
Q6	**Do you have trouble believing that the person who died is truly gone?**					
Q7	**Do you avoid reminders that the person who died is truly gone?**					
Q8	**Do you feel emotional pain related to the death?**					
Q9	**Do you feel that you have trouble re-engaging in life?**					
Q10	**Do you feel emotionally numb or detached from others?**					
Q11	**Do you feel that life is meaningless without the person who died?**					
Q12	**Do you feel alone or lonely without the person who died?**					

Q13. have the symptoms above caused significant impairment in social, occupational, or other important areas of functioning? **Yes / No**

Scoring:
A summary score for the symptom items is calculated by totaling each individual item score from 1 (not at all) to 5 (overwhelmingly) across items 3 through 12. Symptom scores can range from 10 to 50. A score of 30 or greater indicates syndromal-level PGD.